

# **BDS Medium (Gamborg B5 Derivative)**

## **BDS Medium (Gamborg B5 Derivative) in Plant Tissue Culture: Origins, Uses, and Formulation**

### **Origin:**

BDS medium, a derivative of Gamborg's B5 medium, doesn't have a singular, clearly documented "origin" in the way that Murashige and Skoog (MS) medium does. It's more accurate to describe it as a family of media formulations based on modifications of Gamborg's original B5 medium, developed by Ole Gamborg and colleagues in the early to mid-1960s. B5 itself was initially designed for the *in vitro* culture of suspension cultures of various plant cells, including those of soybean and carrot. The goal was to create a defined nutrient medium that supported rapid and consistent cell growth in suspension, a necessary step in the development of plant cell biotechnology at the time. The exact year of its initial publication is debated, with various publications citing slightly different years around the 1960s. The subsequent modifications leading to variations known as "BDS" emerged over time as researchers adapted B5 to specific plant species and culture types, fine-tuning the composition for optimal results. "BDS" isn't a formally standardized acronym like MS; it represents a range of empirically derived modifications of B5.

### **Applications:**

BDS media, in its various modified forms, finds applications across several plant tissue culture techniques. While not universally superior to other media, its versatility makes it suitable for:

- **Callus induction:** BDS is effectively used to initiate callus formation from various explants (e.g., leaf, stem, root segments). Its balanced nutrient composition supports cell division and dedifferentiation.
- **Organogenesis:** Modified BDS media, through careful manipulation of plant growth regulators (PGRs) such as auxins and cytokinins, can promote the development of shoots and roots from callus tissue, enabling plant regeneration.
- **Shoot proliferation:** Specific BDS formulations, typically with higher cytokinin levels, are used for efficient shoot multiplication from nodal segments or pre-existing shoots.
- **Rooting:** Similarly, adjusting the PGR balance in BDS, by increasing auxins and reducing cytokinins, promotes root development in shoots, enabling the successful establishment of plantlets *ex vitro*.

BDS shows particular success with certain plant families, including species from the Solanaceae (e.g., tomatoes, potatoes, tobacco) and certain woody plants though its adaptations vary across the many species. There isn't one definitive large-scale study highlighting BDS's superiority over other media; rather, its effectiveness is demonstrated in numerous smaller publications specific to plant species and experimental conditions.

## Formulation:

The specific formulation of BDS media varies considerably depending on the intended application and the plant species. There is no single "standard" BDS; instead, researchers modify the base B5 medium to optimize specific aspects. A typical base might resemble this (note variations are common even in these components):

| Component   | Concentration (mg/L) | Role                            |
|---|----------------------|---------------------------------|
| $\text{NH}_4\text{NO}_3$                            | 1650                 | Nitrogen source                 |
| $\text{KNO}_3$                                      | 1900                 | Nitrogen and potassium source   |
| $\text{CaCl}_2$                                     | 440                  | Calcium source                  |
| $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$           | 370                  | Magnesium and sulfur source     |
| $\text{KH}_2\text{PO}_4$                            | 170                  | Phosphorus and potassium source |
| $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$           | 27.8                 | Iron source                     |
| $\text{MnSO}_4 \cdot \text{H}_2\text{O}$            | 2.2                  | Manganese source                |
| $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$           | 0.86                 | Zinc source                     |
| KI  | 0.83                 | Iodine source                   |
| $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ | 0.25                 | Molybdenum source               |
| $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$           | 0.025                | Copper source                   |
| $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$           | 0.025                | Cobalt source                   |
| Nicotinic acid                                      | 0.5                  | Vitamin                         |
| Pyridoxine HCl                                      | 0.5                  | Vitamin                         |
| Thiamine HCl  | 0.1                  | Vitamin                         |
| Glycine   | 2                    | Amino acid                      |

| Component               | Concentration (mg/L) | Role   |
|-------------------------|----------------------|--|
| Myo-inositol            | 100                  | Osmolyte and growth regulator                          |
| Sucrose                 | 30000                | Carbon source  |
| Plant growth regulators | Variable             | Auxins (e.g., NAA, 2,4-D), Cytokinins (e.g., BAP, Kin) |

The concentrations of plant growth regulators (PGRs) are critically important and are adjusted depending on the specific objective (callus induction, shoot proliferation, rooting). Different auxin:cytokinin ratios will induce different responses. Other modifications may involve the addition of other vitamins or amino acids, or changes in the concentration of macronutrients based on the plant species and its specific requirements.

### **Conclusion:**

BDS medium, while lacking the formal standardization of MS medium, represents a valuable and adaptable tool in plant tissue culture. Its strengths lie in its efficiency for various plant species including those challenging to cultivate *in vitro*, its relatively simple formulation, and its potential for modification to suit specific needs. Limitations include variations in performance across different plant species requiring specialized adaptation, and it might not be the most suitable medium for all types of culture (e.g. certain orchids or recalcitrant species). Compared to MS medium, BDS often exhibits improved results in specific cases but lacks the widespread documentation and standardization making direct comparison difficult in general. It continues to find relevance in modern plant biotechnology, especially in targeted applications within specific research projects, and

is an example of the ever-evolving flexibility in *in vitro* plant culture techniques.