

Micropropagation of Anthurium – MATSUMOTO 1997



Distribution, Botany, and Importance of *Anthurium*

Within the family Araceae, *Anthurium* is the largest, most morphologically diverse and complex genus, consisting of approximately 1000 species. Native to Central and South America, members of *Anthurium* are found at elevations ranging from sea level to 3000m, most commonly in cloud forests at 1500m (Croat 1986). Plants of this herbaceous perennial monocot are terrestrial or epiphytic. Typical of the amids is the spadix, consisting of a multitude of unobtrusive true flowers supported by a fleshy axil. The protogynous nature of the bisexual flowers in *Anthurium* favors cross-pollination. The commercial flower is a combination of the spadix and a colorful modified leaf, termed spathe. Attractive foliage of some species makes anthuriums also suitable for leaf harvest and cultivation as a potted plant.

Commercial production has focused on plants derived from two major species, *Anthurium andraeanum* Linden ex. Andre and *A. scherzerianum* Schott. (Fig. 1A,B). The majority of the plants used in the cut flower industry are thought to be hybrids of *A. andraeanum* and other species (Madison 1980), and will be referred to as *A. andraeanum* Hort. Main production areas are Hawaii and The Netherlands, with additional production in

other tropical and subtropical regions. The 1991, combined Dutch auctions ranked anthurium 14th of all cut flower sales, with over 20 million stems sold for approximately

\$21.5 million (International Floriculture Quarterly Report 1992). Estimates for Dutch auction anthurium sales in 1993 are approximately 37 million stems (International Floriculture Quarterly Report 1994). In Hawaii, anthurium is one of the top cut flowers, with a 1993 farmgate value of sales of \$7.5 million for 10.6 million stems sold (Hawaii Agricultural Statistics Service 1994). *A. scherzerianum* is sold as a flowering potted plant, with main production areas located in Europe. Global production of anthurium hybrids as potted plants has recently increased.

1 Department of Horticulture, University of Hawaii at Manoa, 3190 Maile Way, St. John 102, Honolulu, Hawaii 96822-2279, USA

² Present address: Department of Horticulture, Purdue University, 1165 Horticultural Building, West Lafayette, Indiana 47906-1165, USA

Biotechnology in Agriculture and Forestry, Vol. 40

High-Tech and Micropropagation VI (ed. by Y.P.S. Bajaj)

© Springer-Verlag Berlin Heidelberg 1997

Micropropagation of *Anthurium* 15



Fig. 1. A *Anthurium andraeanum* Hort., cut flower cultivar

Nitta. **B** *Anthurium scherzerianum*

used in flowering potted plant production. (Photos courtesy H. Kamemoto)

Common Propagation Practices and Need for Micropropagation

Conventional propagation relies upon divisions, cuttings, and in vitro methods. Seeds are less commonly used for propagation, as they may produce heterogeneous populations varying in flower color, size, and form.

Division relies on lateral shoots arising from the basal stem portion of the anthurium plant. Some cultivars produce lateral shoots easily while others produce very few. Time until first harvestable flower depends upon initial size of the division, but is generally in terms of months rather than years. Plant growth regulators have been used to stimulate lateral shoot development

within 4 to 6 months after application. A foliar spray of 1000mg/l benzyladenine (BA) applied to intact Ozaki plants resulted in 3.6 lateral shoots per plant, with zero lateral shoots formed on unsprayed plants (Higaki and Rasmussen 1979). Removal of the apical portion of juvenile Mauna Kea plants followed by a 500mg/l GA₃ spray increased shoot production from 3.3 shoots (without spray) to an average of 5.8 shoots per sprayed plant (Imamura and Higaki 1988).

Top cuttings, consisting of the uppermost stem with two or three leaves, are removed from plants and rooted in a well-aerated medium. Roots develop within 2 to 3 weeks, with the first flower produced in approximately 6 months. Removal of the top cutting stimulates development of lateral shoots on the mother plant. Basal cuttings, consisting of one or two leafless nodal sections and placed horizontally on medium, produce plants from each node. Although more plants may be

generated by basal, rather than top, cuttings, plants take longer to develop and often require 2 to 3 years to reach full production.

In vitro propagation is another method commonly used in anthurium propagation. Plants may be obtained directly from excised apical and lateral buds or indirectly through the differentiation of callus induced from leaf, spathe, and spadix explants.

Rapid clonal propagation is an important use of anthurium tissue culture. In The Netherlands, anthuriums are almost exclusively propagated by culture in vitro (van Doesburg 1991). As recalcitrant genotypes continue to hinder micropropagation of some cultivars, additional in vitro studies should be undertaken.

Disease elimination should be addressed in the propagation of anthuriums. The bacterial blight caused by *Xanthomonas campestris* pv. *dieffenbachiae* has been a problem in Hawaii for a decade, and occurs in other anthurium-producing countries such as the Philippines, Jamaica, Tahiti, and Trinidad. *Xanthomonas campestris* pv. *dieffenbachiae* may be present in callus and stage II plantlets without visible symptoms, and will not cause medium turbidity in MS-based media that lack coconut water (Norman and Alvarez 1994). Efforts are underway to develop an indexing and certification program for micropropagated anthuriums (Tanabe et al. 1992; Fernandez et al. 1992).

Due to the long breeding cycle of anthuriums, in which the development of a new cultivar may take 8 to 10 years, genetic engineering as a viable breeding aid is currently being investigated at the University of Hawaii for resistance to bacterial blight (Kuehnle et al. 1992b).

Review of In Vitro Studies

Early work on anthurium tissue culture described callus proliferation and subsequent plantlet formation from seeds and young leaf tissue of mature anthurium plants (Pierik et al. 1974). Following this pioneering work, many studies have been conducted for several species and hybrids (reviewed by Geier 1990; Table 1). Different culture media containing modifications of

Table 1. Summary of micropropagation studies in *Anthurium*. (See also Geier 1990)”

Anthurium species/hybrids	Explant material	Observations/remarks	Reference
<i>A. andraeanum</i>	In vitro plant	Acclimatization of rooted plants	Imamura and Higaki (1981)
<i>A. scherzerianum</i>	Spadix fragments	Ploidy variation in callus and regenerated plants	Geier (1982,1988)
<i>A. andraeanum</i>	Seed	Multiple plant formation	Tanabe et al. (1989)
		from seeds germinated in	
		vitro	
<i>A. andraeanum</i>	Lamina	Callus and plant regeneration on medium with BA and/or 2,4-D	Lightbourn and Devi Prasad (1990)
<i>A. andraeanum</i>	In vitro plant	Certification and indexing	Fernandez et al. (1992)

		procedure for detection of	Tanabe et al. (1992)
		<i>Xanthomonas campestris</i> pv. <i>diffenbachiae</i>	
<i>A. andraeanum</i>	Etiolated shoot	Protoplast obtained but sustained division not observed	Kuehnle and Nan (1991)
<i>A. andraeanum</i>	Lamina and petiole	Shoot and roots from	Kuehnle and Sugii
		caulogenic callus, genotype effect on regeneration	(1991)
<i>A. andraeanum</i>	Lamina and spadix	Spadix explants produced more callus formation and plant regeneration than	Singh and Sangama (1991)
		lamina, with greater ploidy	
		uniformity; some	
		aneuploids were obtained	
<i>A. andraeanum</i>	In vitro plant	Stage II and stage III plant acclimatization	Tanabe (1991)
<i>A. andraeanum</i>	Bud	Effect of genotype on time to first leaf formation	Tanabe et al. (1991)

<i>A. andraeanum</i> <i>A. lindenianum</i> <i>A. amnicola</i> <i>A. kamemotoanum</i>	In vitro lamina	Somatic embryos obtained and produced single plants or multiple plant clumps	Kuehnle et al. (1992)
<i>A. scherzerianum</i>	Lamina and petiole	Caulogenic callus and plant regeneration	Liu and Xu (1992)
<i>A. andraeanum</i>	Bud	Surface disinfestation of buds	Tanabe and Matsumoto (1992)
<i>A. cubense</i>	Lamina and petiole	Examined leaf stage, and medium NH₄NO₃ and 2,4-D on callus formation	Warner et al. (1993)
<i>A. andraeanum</i>	Lamina, petiole, callus and plantlets	<i>Xanthomonas</i> <i>campestris</i> pv. <i>diffenbachiae</i> may be present in callus and stage II shoots without visible symptoms or medium turbidity	Norman and Alvarez (1994)

·' References summarized in Geier (1990) are not included.

Nitsch (1969) or MS (Murashige and Skoog 1962) basal salts, sugars, other organic components, and growth regulators were described for proliferation and plant regeneration from a variety of tissues. Axillary bud culture (Kunisaki 1980) proved to be another effective micropropagation method. Effects of explant and genotype on regeneration and genetic stability of regenerated plants have also been studied.

Recent studies describe callus, somatic embryogenesis, and protoplast culture. Caulogenic callus was induced from leaf and petiole segments of *A. scherzerianum*, multiple shoots were obtained, and rooting was induced (Liu and Xu 1992). *A.*

andraeanum Hort. callus and shoot regeneration was reported by Lightbourn and Devi Prasad (1990), Kuehnle and Sugii (1991), and Singh (1991). Trends in these studies were similar to those reviewed in Geier (1990), where callus is induced on a modified Nitsch (1969) or Pierik et al. (1974) medium supplemented with only 2,4-dichlorophenoxyacetic acid (2,4-D) or a combination of 2,4-D and BA. Plant regeneration was achieved on medium with BA or devoid of growth regulators. A method for somatic embryo induction and conversion to plants was developed for *A. andraeanum* hybrids using in vitro lamina as explant sources (Kuehnle et al. 1992a). Preliminary work on anthurium protoplast isolation and culture was presented by Kuehnle and Nan (1991).

Micropropagation

1. Establishment of Axenic Cultures

Disinfestation of anthurium tissues can be problematic, with the exception of seeds. Contamination rates usually range from 10 to 20% among leaf explants, 75% in spadix sections (Geier 1990), and 33 to 87% for excised axillary buds (Kunisaki 1980). Due to the protective layers of berry flesh and seed coat, disinfectant may be used at higher concentrations or longer exposure times with fruit and seeds. Slow-growing contaminants are not unusual in anthurium tissues and may appear after the first month of culture. Successfully disinfested explants should show minimal discoloration.

1. Seed

Seeds may be disinfested by first soaking the harvested berry in 3% sodium hypochlorite (NaOCl; e.g., 57.7% Clorox) for 15min, followed by soaking excised seeds in 1 % NaOCl for 20min (Pierik et al. 1974). Each soak is accompanied by a 30-min rinse with several volumes of sterile water. The seed coat is then removed, and the explant, consisting of the embryo and endosperm, is cultured on appropriate medium (Pierik et al.

1974). Calcium hypochlorite can be substituted for sodium hypochlorite (Rosario and Lapitan

1981). Successful disinfestation has also been achieved by one soak of excised seed in 2.6% NaOCl for 5min (Zens and Zimmer 1988) or LD disinfectant (Alcide Corp., Norwalk, Connecticut) at 1 part activator: 1 part base: 10 parts water for 30min (Tanabe et al. 1989). Presence of a gelatinous or sticky substance often hinders the handling of the anthurium seeds with standard tissue culture tools. Although it is not usually present in seeds disinfested with NaOCl, removal of this substance is possible with a 13% sodium carbamate solution (Maurer and Brandes 1979). Seeds or excised embryos germinate within 4 weeks, with proliferation of callus usually occurring within 12 to 16 weeks (Rosario and Lapitan 1981).

1. Leaf, Spathe, and Spadix

Lamina, petiole, and spathe sections are generally disinfested by an initial dip in 70 to 95% alcohol, followed by a 10- to 30-min soak in 1.5 to 3% NaOCl. An alternative method uses a 5-min soak in 0.1% mercuric chloride solution with

0.25 ml/l Tween 20 in place of NaOCl (Eapen and Rao 1985). Similar methods are used for spadix explants, in which first the spathe surrounding the young spadix is disinfested, followed by disinfestation of the spadix proper (Geier 1982).

Unprotected or screenhouse cultivation of plants in subtropical and tropical areas is conducive to high contamination rates for field-grown material. In Jamaica, use of a 70% alcohol dip for 45s and a 1.25% NaOCl soak for 15min resulted in up to 70% contamination of leaf explants. Contamination was reduced to 10% by a presterilization soak in the fungicide Benlate (Dupont) (Lightbourn and Devi Prasad 1990). Axenic leaf blade and petiole cultures have been obtained using an initial 10-min soak in 0.14% Phytan 20 (Maril Products Inc., Tustin, California) followed by

consecutive soaks of 30min in 0.53%, then 0.27%, NaOCl with one drop Tween 20 in 100ml (Kuehnle and Sugii 1991); contamination rates of 5% are routinely achieved (T. Matsumoto, unpubl.).

While leaf callus produced from plants infected by *Xanthomonas campestris* pv. *dieffenbachiae* was found to be axenic, it should be noted that this bacterial pathogen can be harbored asymptotically in inoculated callus and shoots for 4 months to 1 year (Norman and Alvarez 1994).

1. Axillary Buds

Initial reports record a contamination rate of 33% using two soaks for 20 and 45min in 0.53 and 0.27% NaOCl solution with Tween 20, respectively, with removal of bud scales (Kunisaki 1980). Reduction of exposure time to the disinfectant is possible through use of LD and Exspor disinfectants supplemented with 35% isopropyl alcohol (Alcide Corp., Norwalk, Connecticut; Tanabe and Matsumoto 1992).

Methods for Culture Initiation and Plant Multiplication

Enhanced axillary branching from microcuttings of in vitro shoots is used to multiply plants to the quantities desired. The initial shoots are obtained from a variety of sources, namely callus cultures, axillary buds, and somatic embryos. Several factors affect the success of anthurium micropropagation, and some aberrations have been described among regenerated plants.

3.2.1 Callus

Callusing is used to initiate [in vitro cultures](#) of field material from which shoots for subsequent multiplication are derived (Fig. 2). This method is commonly used in The Netherlands and numerous reports have been published.

Organogenic callus and plant regeneration have been achieved using seeds, embryos, and explant material of leaf lamina, petiole, spadix, spathe, and etiolated shoots (Geier 1990; Lightbourn and Devi Prasad 1990; Kuehnle and Sugii 1991; Liu and Xu 1992). In general, callus induction and proliferation are favored under dark conditions by addition of an auxin, usually 2,4-D, and a cytokinin, usually BA, to solid or liquid medium. Shoot proliferation from callus is stimulated with the removal of auxin from the medium, reduction of ammoniacal nitrogen, and increased light. Cytokinins such as 2-isopentenyladenine (2iP), BA, or kinetin may, in some cases, be required for shoot formation.

The soft tissue of newly unfolded leaves is successfully used for laminar explants. Explants of fully expanded leaves should include a major vein with




Fig. 2. Callus initiated from Kozohara lamina and Nitta petiole sections on modified Pierik medium with $0.36\mu\text{M}$ 2,4-D and $4.4\mu\text{M}$ BA with 0.18% Gelritc in the dark


Micropropagation of *Anthurium* 21

vascular tissue for improved proliferation (Finnie and van Staden 1986). It is suggested that young, un lignified leaves about one-half to two-thirds of the final length are most useful for *A. scherzerianum* (Geier 1990). Optimum regeneration in *A. andraeanum* occurs if leaves are harvested 1 day after they are fully expanded. Lamina sections show signs of proliferation as early as 2 to 4 weeks (Finnie and van Staden 1986) up to 12 to 16 weeks (Lightbourn and Devi Prasad 1990).

1. Axillary Bud

Direct shoot formation from excised buds was reported by Kunisaki (1980). This method establishes initial shoot cultures without an intervening callus phase, and thus may reduce the possibility of somaclonal variation and abnormal plant recovery, but at the expense of rapid propagation. According to Kunisaki (1980) and later adaptations (J. Kunisaki, pers. comm.) five to ten lateral buds are obtained from the stem of an anthurium plant (Fig. 3A), surface sterilized, and trimmed to 2 mm at the base. Shoot formation is encour-

A



B

Fig. 3A,B. Axillary bud culture. **A** Stem section of mother plant, stripped of leaf sheaths to expose lateral buds. **B** Multiple shoot formation from stem sections of plants grown from axillary buds on modified half-strength MS medium with 2% sucrose and $0.89\mu\text{MBA}$. (Photos courtesy J. Kunisaki)

aged in a liquid modified MS medium consisting of MS salts at

3/8 strength, 15% coconut water, and 2% sucrose. After a single elongated shoot develops from each bud, usually within 12 to 18 months, a top cutting consisting of the apex and two or more leaves is cultured on a filter paper bridge in similar medium. For multiple shoot formation, basal portions of the remaining stem are placed on solid medium, or on a shaker in liquid medium, supplemented with 0.2 mg/l BA for a maximum of 2 months (Fig. 3B). Top cuttings taken from the multiple shoots are placed on medium used for initial shoot formation and solidified with 0.18% Gelrite for shoot growth and root formation. The basal explants are subcultured again to solid medium lacking BA for additional shoot formation. Once shoots form, top cuttings are taken once again and the remaining bases discarded.

1. *Somatic Embryogenesis*

Somatic embryogenesis and subsequent plant regeneration have been achieved in *A. andraeanum* hybrids. Whole lamina explants were harvested from in vitro-grown plants and plated on a modified half-strength MS medium with 2% sucrose, 1% glucose supplemented with 1 to 4mg/12,4-D, and 0.33 to 1 mg/l kinetin. Induction of embryos (Fig. 4) and proliferation of secondary embryos occurred under darkness. Conversion and maturation occurred on the same basal medium plus 2% sucrose, 0.2mg/l BA, and 0.18% Gelrite under a 16-h photoperiod (Kuehnle et al. 1992a). These plantlets can serve as a source of nodal microcuttings for subsequent multiplication.

In other reports, somatic embryogenesis in spadix callus of *Anthurium scherzerianum* was induced by lowering ammonium nitrate to 1.25 mM




Fig. 4. Somatic embryos of *Nitta* cultured in the dark on modified half-strength MS medium with 2% sucrose, 1% glucose, 18 μ M 2,4-D, and 2.3 μ M kinetin

Micropropagation of *Anthurium* 23

NH_4NO_3 in a Nitsch medium with 4.44 μ M BA and 0.45 μ M 2,4-D (Geier 1982). Somatic embryos germinated into bipolar structures and multiple plantlet clumps. However, these somatic embryos occurred sporadically (Geier I 990). Petioles from in vitro-grown plants of Lady Jane (*A. andraeanum* X *A. antioquiense*) plated on a modified MS medium with 2 μ M 6-benzylaminopurine (BAP), 2 μ M zeatin, and 1 μ M 2iP, also appeared to produce somatic embryos (F.J. Novak, pers. comm.). Cultures were kept in the dark for 2 to 3 months at 28 °C and maintained on the same medium, either solid or liquid. Shoots were regenerated on a medium containing 0.5 μ M IBA (indolebutyric acid).

1. Influence of Genotype and Selection of Explant

Genotype plays an important role in the multiplication and regeneration of anthuriums (Pierik et al. 1974). As with other crops, examples of variation among genotypes exist for anthurium callus formation (Geier 1990; Light bourn and Devi Prasad 1990; Kuehnle and Sugii 1991; Liu and Xu 1992) and somatic embryogenesis (Kuehnle et al. 1992a). Differences in cultivar responses also apply to enhanced axillary branching, with the period of first leaf emergence ranging from 13 weeks for Hawaiian Butterfly to 38 weeks for Fuji Pink (Tanabe et al. 1991).

Selection of the explant is often dependent upon the material available and the objective of study. Limited number of axillary buds are available per individual mother plant, and recovery of the mother plant following removal of a large stem section for excision of axillary buds may take over one year. In contrast, greater amounts of propagative material are generally available for other methods, and removal of a leaf, spathe, or spadix will not cause substantial injury to a mother plant.

For clonal propagation, the use of seeds is highly discouraged due to the high variability in offspring from cross-pollinations. However, seed culture has been proposed as a breeding aid applicable to limited number of crosses or those with few resulting seeds (Tanabe et al. 1989). Clones are produced in vitro by promotion of multiple shoot formation from germinating seeds. Clones of each seed are evaluated by the breeder in the field; corresponding clones are retained in vitro. If and when the selection of a promising cultivar is made, propagules are already available in vitro for subsequent multiplication.

1. Rooting

Shoots will spontaneously root in culture under light conditions following removal or depletion of growth regulators, notably cytokinins, in the nutrient medium. If rooting does not readily occur, medium salts may be reduced to half the original strength and the solidifying agent may be reduced or eliminated. Some propagators include charcoal as a darkening agent in the rooting medium (N. van der Knaap, pers. comm.).

1. Acclimatization

Anthurium microcuttings rooted in vitro (termed stage III) prior to transfer to the greenhouse are generally hardier than unrooted propagules (stage II), but may be less cost-effective for the micropropagator or grower. Retail price per stage II microcutting is currently 20% less than stage III material for major labs in the USA. Success of acclimatization at stage II is often dependent on the particular cultivar. Many labs also sell acclimatized plants (Stage IV). These often ensure a better unit price per plant for the laboratory and increase survivability of plants for the buyer. Plants at any of these stages are susceptible to bacterial blight, rain and slug damage, and salt burn from use of undiluted complete fertilizer.

Stage II and stage III microcuttings should be carefully removed from the culture vessels and all remaining agar rinsed off. Stage II shoots preferably should contain the apex plus two or three leaves prior to transplantation ex vitro. Microcuttings are placed into a sterile, premoistened medium such as Oasis Root Cube, rockwool, or vermiculite/perlite combinations. Plants are kept in high humidity under 80% shade or 1500 to 2000 foot candles for approximately 2 months (Tanabe 1991). Stage III plantlets may be treated with a fungicide as a preplant soak or postplant drench (Imamura and Higaki 1981). Plants with intact roots are transplanted into a variety of media including Oasis or rockwool plugs, shredded tree fern fiber, or mixes of soil amendments such as

composted shredded bark: no. 2 perlite (3:1 mix) under 80% shade with high humidity.

In large-scale commercial productions, misting or fogging systems are often employed to provide high humidity conditions conducive for further plant growth. Other methods include use of humidity tents for multiple flats (Fig. SA) or clear plastic domes for individual flats (Fig. SB). Plants are monitored and watered to prevent desiccation. The plastic cover should be removed in incremental stages, to gradually reduce the humidity (Tanabe 1991). Plants should continue to be protected from rain until large enough for transplant into beds or pots (Higaki et al. 1994).

1. Somaclonal Variation

Aberrations are occasionally observed among micropropagated plants. Although ploidy variation was found in callus of *A. scherzerianum*, carry-over to the regenerated plant was rare, based on stoma length, chromosome counts, and cytophotometry (Geier 1988, 1990).

Another source of variation may relate to the proliferation method used. Somaclonal variation was documented for plants obtained both from year old callus pieces and from plants initiated by axillary bud culture and micropropagated long-term by enhanced axillary branching (Kuehnle and Sugii 1992; Kuehnle and Kuanprasert, unpubl.). Using 3 diverse genotypes, approximately 520 plants micropropagated via long-term callus culture (12 to 13 months) and 120 plants micropropagated via nodal cuttings, over a 3- to 4- year period, were grown to maturity and evaluated. Plants were grouped into

Micropropagation of *Anthurium* 25



Fig. SA,B. Acclimatization: **A** Stage III anthuriums

transplanted ex vitro into trays containing Oasis rooting cubes and covered with plastic hoods. **B** Humidity tent used to cover mass plantings of anthurium

those arising from the same callus piece immediately prior to shoot regeneration, or those arising from the same nodal cuttings in the last subcultures prior to planting in the greenhouse.

Both sets of plants experienced stunting or shortening of internodes for all three genotypes. Shortened internodes occurred among 19 to 30% of the callus groups, but infrequently within a group, so that as few as 4% of the total

plants from a genotype may have had reduced stature. Plants from nodal cuttings showed a higher rate of stunting (25 to 43% of related groups), but no other aberrations. Among callus-derived plants, leaf shape abnormalities, leaf variegation or ploidy changes were detected in 0.8 to 2% of the total plants, corresponding to 3 to 8% of the original callus clumps put on plant regeneration medium; frequency varied among genotypes.

Based on these findings, it is suggested that callus proliferation be restricted to a few months prior to plant regeneration and that new stock plants for enhanced axillary branching, derived from callus or from axillary buds, be re-initiated at regular intervals of less than 3 years; every 6 to 12 months is practiced in The Netherlands.

1. Commercial Aspects (Table 2)

Among commercial tissue culture laboratories in 14 Western European countries surveyed in 1988, *Anthurium* ranked 14th among the most frequently propagated genera, with three countries producing more than 100000 anthurium plants each (Pierik 1991). The Netherlands, which accounts for 29% of the West European production, micropropagated over 0.5 million plants of *A. andraeanum* in 1988, placing *A. andraeanum* second

after *Gerbera* in quantity of plants produced for the cut flower industry. For the potted plant market, *A. scherzerianum* ranked fourth, after *Nephrolepis*, *Saintpaulia*, and *Ficus*, with 1.7 million plants produced in vitro in The Netherlands (Pierik 1991). Figures for the quantity of plants micropropagated from 1988 to 1990 have been compiled and are shown in Table 2 (R.L.M. Pierik, pers. comm.). In The Netherlands, initial multiplication relies largely upon callus culture.

To estimate the extent of *Anthurium* micropropagation in the USA, a survey was sent to major commercial tissue culture laboratories in Hawaii and the continental USA. Of seven laboratories currently producing over 10000 anthuriums per year, an estimated 3.8 million plants were micropropagated in 1993, with three of these labs producing 90% of the supply with over 1 million plants each. The majority of plants were for use in cut flower and flowering potted plant production; fewer were produced for the foliage industry. A variety of *A. andraeanum* Hort., *A. scherzerianum*, and hybrids with *A. amnicola* and *A. antioquiense* are produced for the flowering potted plant market. All seven laboratories rely primarily on axillary and apical bud culture

Table 2. Quantity of micropropagated anthurium plants in The Netherlands, 1988 to 1990. (Courtesy R.L.M. Picrik, Agricultural University, Wageningen, The Netherlands)

	1988	1989	1990
<i>A. andraeanum</i>	509300	919000	1516500
<i>A. scherzerianum</i>	1684624	2454400	4157750
Total plants	2193924	3373400	5674250

for culture establishment, followed by axillary branching for multiplication; two of seven (29%) labs use callus or somatic embryogenesis as a supplemental form of culture establishment. The vast majority of plants sold are derived

from initial bud culture. Micropropagated anthurium plants are sold by 71% of the laboratories as stage III or stage IV; 29% of the labs also offer stage II microcuttings. In terms of total plants sold, stages III and IV are slightly more popular, with an estimated 35% stage III, 37% stage IV, and 28% stage II sold by the seven laboratories.

Summary and Conclusions

Methods for fast clonal propagation are essential to fulfill growers' demands for anthurium plants worldwide. Heterogeneous seed progeny and slow methods of field propagation are unsuitable for generating the large quantities needed. Efficiency of callus culture has been greatly improved for many genotypes since this system was first described for anthurium in 1974 by Pierik et al. Alternative methods, such as axillary bud culture and somatic embryogenesis, have also been developed. Currently, all three methods are commercially used for micropropagation of cut flower and potted plant anthuriums. Slow response of certain genotypes is still a problem despite modifications to the culture medium. Mass propagation linked with disease-indexing programs are currently being investigated, and will be beneficial to production areas plagued with the bacterial blight. In the future, in vitro regeneration of transgenic plants might play an important role in breeding for disease resistance and other desirable qualitative traits.

1. Protocol

Axillary Bud

1. Excise buds with a 1-cm base from a stem which has been washed with soap, soaked in 0.53% NaOCl for 5 min, and air-dried for 2-3 days.
2. Soak in 0.53% NaOCl for 30 min, rinse, then remove two to three leaf sheaths and cut to 2 mm.

3. Soak again in disinfestation solution for 5 min and rinse in sterile water.
4. Place bud in liquid half-strength MS medium with 2% sucrose, 15% coconut water in culture tube on rotary drum at 0.2 rpm, under light, until shoot elongates with leaves.
5. Use basal stem cutting for enhanced axillary branching on half-strength MS medium with 2% sucrose and 0.2mg/l BA under light; shoots are rooted on hormone-free medium in light.

Callus

1. Surface sterilize petiole and lamina sections in 0.14% Phytan 20 for 10min, 0.53% NaOCl for 30min, and 0.26% NaOCl for 30min, then rinse in sterile water.
2. Plate on solid modified MS medium containing 1 mg/l BA and 0.08-0.1 mg/l 2,4-D in dark.
3. After several months, explants with callus are moved into light for shoot proliferation on BA medium.
4. Cuttings are taken for enhanced axillary branching and rooting, as in 5.1.5, above.

References

Croat T (1986) The distribution of *Anthurium* (Araceae) in Mexico, Middle America and Panama.

Selbyana 9: 94-99

Eapen S, Rao PS (1985) Regeneration of plants from callus cultures of *Anthurium patulum*. Curr Sci 54: 284-286

Fernandez JA, Tanabe M, Crane S, Wolff W, Moriyasu P (1992) *Anthurium* in vitro triple indexing II. In: Delate KM, Tome CHM (eds) Proc 5th Hawaii Anthurium Industry Conf. Hawaii Institute of Tropical Agriculture and Human Resources, HIT AHR 02.02.94, University of Hawaii, Honolulu, p 19

Finnie JF, van Staden J (1986) In vitro culture of *Anthurium andraeanum*. S Afr J Bot 52: 343-346
Geier T (1982) Morphogenesis and plant regeneration from spadix fragments of *Anthurium scherzerianum* cultivated in vitro. In: Fujiwara A (ed) Plant tissue culture 1982. Maruzen,

Tokyo, pp 137-138

Geier T (1988) Ploidy variation in callus and regenerated plants of *Anthurium scherzerianum*

Schott. Acta Hort 226: 293-298

Geier T (1990) Anthurium. In: Ammirato PV, Evans DA, Sharp WR, Bajaj YPS (eds) Handbook of plant cell and tissue culture, vol 5. Ornamental species. McGraw-Hill, New York, pp 228- 252

Hawaii Agricultural Statistics Service (1994) Hawaii flowers and nursery products, annual summary. Hawaii Dep Agric, US Dep Agric, Honolulu, Hawaii

Higaki T, Rasmussen HP (1979) Chemical induction of adventitious shoots in anthurium.

HortScience 14: 64-65

Higaki T, Lichty JS, Moniz D (1994) Anthurium culture in Hawaii. HIT AHR Research Extension Series 152, University of Hawaii, Honolulu

Imamura J, Higaki T (1981) Transfer of tissue-cultured anthurium plantlets from flask to natural conditions. Hortic Digest 61: 1-2

Imamura J, Higaki T (1988) Effect of GA₃ and BA on lateral shoot production on *Anthurium*.

HortScience 23: 353-354

International Floriculture Quarterly Report (1992) Cut flower 1991 rankings, vol 3. Pathfast Publishing, Essex, pp 56-62

International Floriculture Quarterly Report (1994) Cut flower and pot plant prices through the Dutch auctions, vol 4. Pathfast Publishing, Essex, pp 63, 77-78

Kuehnle AR, Nan GL (1991) Isolation and culture of protoplasts from two tropical monocots,

Anthurium and *Dendrobium* orchid. *Physiol Plant* 82: A7

Kuehnle AR, Sugii N (1991) Callus induction and plantlet regeneration of Hawaiian anthuriums.

HortScience 26: 919-921

Kuehnle AR, Sugii N (1992) Update on somatic embryogenesis research. In: Delate KM, Tome CHM (eds) Proc 5th Hawaii Anthurium Industry Conf. Hawaii Institute of Tropical Agriculture and Human Resources, HIT AHR 02.02.94, University of Hawaii, Honolulu, pp 15-16

Kuehnle AR, Chen FC, Sugii N (1992a) Somatic embryogenesis and plant regeneration in *Anthurium andraeanum* hybrids. *Plant Cell Rep* 11: 438-442

Kuehnle AR, Chen FC, Sugii N, Jaynes J, Norman D, Alvarez A (1992b) Engineering blight resistance in *Anthurium*: a progress report. In: Delate KM, Tome CHM (eds) Proc 5th Hawaii Anthurium Industry Conf. Hawaii Institute of Tropical Agriculture and Human Resources, HIT AHR 02.02.94, University of Hawaii, Honolulu, pp 17-18

Kunisaki JT (1980) In vitro propagation of *Anthurium andreanum* Lind. *HortScience* 15: 508-509 Lightbourn GJ, Devi Prasad PV (1990) In vitro techniques for rapid multiplication of four

varieties of *Anthurium andraeanum* in Jamaica. *Proc Interam Soc Trop Hort* 34: 3-5

Liu CM, Xu ZH (1992) An efficient procedure for micropropagation of *Anthurium scherzerianum*

Schott (flamingo flower). Chin J Bot 4: 49-55

Madison M (1980) Aroid profile no. 6: *Anthurium andreanum*. Aroideana 3: 58-60

Maurer M, Brandes S (1979) Die Extraktion der Samen von *Anthurium scherzerianum*-Hybriden.

Gartenbauwissenschaft 44: 71-73

Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiol Plant 15: 473-497

Nitsch JP (1969) Experimental androgenesis in *Nicotiana*. Phytomorphology 19: 389-404 Norman D, Alvarez A (1994) Latent infections of in vitro anthurium caused by *Xanthomonas*

campestris pv. *dieffenbachiae*. Plant Cell Tissue Organ Cult 39: 55-61

Pierik RLM (1991) Commercial micropropagation in Western Europe and Israel. In: Debergh PC, Zimmerman RH (eds) Micropropagation. Kluwer, Dordrecht, pp 155-165

Pierik RLM, Steegmans HHM, van der Meys JAJ (1974) Plantlet formation in callus tissues of

Anthurium andraeanum Lind. Sci Hortic 2: 193-198

Rosario TL, Lapitan LA (1981) Callus and plantlet formation in *Anthurium andraeanum* Lind.

Phillipp Agric 64: 197-202

Singh F, Sangama (1991) Micropropagation and plant conformity in *Anthurium andraeanum*. In: Prakash J, Pierik RLM (eds) Horticulture – new technologies and applications. Kluwer, Dordrecht, pp 201-204

Tanabe M (1991) In vitro plant acclimatization. In: Leonhardt

K, Evans D, Halloran J (eds) Hawaii tropical cut flower industry conference: growing in the 90's. Hawaii Institute of Tropical Agriculture and Human Resources, HITAHR Res Ext Series 124, University of Hawaii, Honolulu, pp 100-101

Tanabe M, Matsumoto T (1992) Anthurium explant surface disinfestation. In: Delate KM, Tome CHM (eds) Proc 5th Hawaii Anthurium Industry Conf. Hawaii Institute of Tropical Agriculture and Human Resources, HITAHR 02.02.94, University of Hawaii, Honolulu, pp 10-11

Tanabe M, Keolanui R, Campbell J (1989) Anthurium seed tissue culture. In: Fernandez JA, Nishijima WT (eds) Proc 2nd Anthurium Blight Conf. Hawaii Institute of Tropical Agriculture and Human Resources, HITAHR 03.10.89, University of Hawaii, Honolulu, pp 52-53

Tanabe M, Arakawa C, Matsumoto T, Tanaka R, English J (1991) Anthurium in vitro propagation. In: Alvarez AM, Deardorff DC, Wadsworth KB (eds) Proc 4th Hawaii Anthurium industry conf. Hawaii Institute of Tropical Agriculture and Human Resources, HITAHR 06.18.91, University of Hawaii, Honolulu, pp 4-6

Tanabe M, Fernandez J, Moriyasu P, Crane S, Wolff W, Liu RW (1992) Anthurium in vitro triple indexing. In: Delate KM, Tome CHM (eds) Proc 5th Hawaii Anthurium Industry Conf. Hawaii Institute of Tropical Agriculture and Human Resources, HITAHR 02.02.94, University of Hawaii, Honolulu, pp 8-9

van Doesburg J (1991) What it takes to be successful: promotion, an essential factor in marketing. In: Leonhardt K, Evans D, Halloran J (eds) Hawaii tropical cut flower industry conference: growing in the 90's. Hawaii Institute of Tropical Agriculture and Human Resources, HITAHR Res Ext Series 124, University of Hawaii, Honolulu, pp 22-31

Warner J, Herrera J, Guevara E (1993) Morphogenesis in vitro de *Anthurium cubense* (Araceae).

Rev Biol Trop 41: 455-460

Zens A, Zimmer K (1988) Development of new cultivars of *Anthurium scherzerianum* Schott through in vitro propagation III. In vitro regeneration from leaf explants of clones. Gartenbauwissenschaft 53: 170-173